

Lidonostrum Bomba-Spray 10%



Nasal, Oral and Dental
Procedures



Oropharyngeal
Procedures



Respiratory Tract
Procedures



Obstetrics and
Gynecology
Procedures



Active Substance
Lidocaine base



Strength | Dosage Form
100 mg/ml Cutaneous spray, solution



Presentation
80 g bottle



Storage conditions
Store below 30° C



Shelf-life
3 Years (in-use: 12 months)



Each spraying applies 10 mg of lidocaine. It is not necessary to dry the site before application.

For specific information on the recommended dose and maximum dose, please refer to the SPC.



To be used in the mucous membranes and provides effective superficial anaesthesia for approximately 10 to 15 minutes.

Anaesthetic action onset usually begins within 1-3 minutes depending on the site where it is applied.

Posology

| AREA | RECOMMENDED DOSE (mg) |
|---|-----------------------|
| Nasal procedures, e.g. puncture of the maxillary sinus. | 20 - 60 |
| Oral and dental procedures, e.g. prior to injection. | 20 - 200 |
| Oropharyngeal procedures, e.g. gastrointestinal endoscopy. | 20 - 200 |
| Respiratory tract procedures, e.g. introduction of instruments and catheters. | 50 - 400 |
| Procedures in the larynx, trachea and bronchi. | 50 - 200 |
| Obstetric procedures, e.g. vaginal delivery, suture of the mucosae and cervical biopsies. | 50 - 200 |

Precautions for use

Dose recommendations must be considered as guidance, the experience of the doctor and the knowledge of the general condition of the patient being essential for the determination of the dose.

Use with caution in patients with wounds or traumatised mucosa in the application site, due to increased systemic absorption.

The use in the oropharynx may interfere with swallowing, thus increasing the danger of aspiration and biting trauma.

Avoid contact with the eyes.

