

Lidonostrum Gele 2%



Respiratory Tract
(nasogastric tubes,
endotracheal intubation,
bronchoscopy)



Digestive Tract
(endoscopy,
procto-rectoscopy)



Urinary System
(catheterization,
cystoscopy)



Active Substance
Lidocaine hydrochloride



Presentation
30 g aluminium tube



20 mg/g
Strength | Dosage Form
20 mg/g Gel



Storage conditions
Do not store above 30° C



Route of Administration
Cutaneous use



3 Years
Shelf-life
3 Years (In-use: 12 months)





In general, the absorption is higher in the bronchial tree, less significant in pharyngeal use and much reduced in endourethral use.



To be used in the anaesthesia of mucosae having a duration of approximately 20 to 30 minutes and a very quick onset, usually up to 5 minutes.

Posology

RESPIRATORY TRACT

Bronchoscopy and Endotraqueal intubation

- Bronchoscopy: 10-20 ml gel.
- Endotracheal intubation: 2 ml of gel in the surface of the catheter. Do not allow the gel to go inside the catheter.

DIGESTIVE TRACT

Endoscopy

- Instillation of 10-20 ml.
- Gel can be applied in the instrument as lubricant.
- When combined with other lidocaine-containing products - total dose must not exceed 400 mg.

URINARY SYSTEM

Urethral anaesthesia - surface anaesthesia

- Male: 20 ml of gel (400mg), maximum of up to 40 ml of gel divided into 3-4 applications.
- Female: 5-10 ml of gel in small amounts.

Precautions for use

Elderly or debilitated patients, children aged ≥ 12 years old, patients with acute diseases or infections must receive doses adapted to their age, weight and physical condition.

Lidocaine is considered porphyrinogenic in animals, and thus, not considered safe in patients with acute porphyria.

Contraindicated in the very rare cases of patients with hypersensitivity to local anaesthetics of the amide type or to any other components of the gel.

This medicinal product contains parabens.

