



GRUPO

**Caring** for science.  
**Caring** for you.

# PRODUCT LIST



PAIN & FEVER  
COUGH & THROAT  
ANTI-INFLAMMATORY  
GASTROENTEROLOGY  
ANTIVIRAL  
ANTIBACTERIAL  
CARDIOLOGY  
CENTRAL NERVOUS SYSTEM  
GLUCOCORTICOID  
WARTS TREATMENT  
CALCIUM FOLINATE  
ANTIHISTAMINIC  
ANTIDISLIPIDEMIC  
BENIGN PROSTATIC HYPERPLASIA  
VITAMINS & MINERALS  
HAIR LOSS  
HAIR CARE  
ANTI-SEBORRHEIC  
ACNE  
CICATRIZATION  
DRY & PSORIATIC SKIN  
BABY CARE  
BATH & HYDRATION  
ANTI-AGING  
ANIMAL HEALTH

# index

## PRODUCT LIST



**MEDINFAR** is a fully integrated Pharmaceutical Company with headquarters in Lisbon, **Portugal**. Founded in 1970, Medinfar is focused in R&D, Production, Distribution, Marketing and Sales, being currently the N°1 Portuguese Company in Consumer Health and Dermatology. In Portugal, besides our own brands, we market licensed products in partnerships with top international pharmaceutical companies.

**Medinfar markets a wide range of products:** Prescription medicines, Consumer Health, Dermocosmetics, Food Supplements and Veterinary products, **in more than 70 countries.**

Through our **R&D - GMP center** and our **EU-GMP manufacturing** site Medinfar Manufacturing, we offer tailor made turnkey solutions, from galenic development to contract manufacturing.

We offer

**LICENSING OUT**  
**CONTRACT MANUFACTURING**  
**CONTRACT DEVELOPMENT**

# human health

## LABORATÓRIO MEDINFAR

Marketing and Sales of Prescription  
Branded Products

## MEDINFAR CONSUMER HEALTH

Marketing and Sales of OTC and Other  
Consumer Health Products

## GP GENÉRICOS PORTUGUESES

Marketing and Sales of Prescription  
Generic Products

# manufacturing

## MEDINFAR MANUFACTURING

EU • Manufacturing Plant

# animal health

## MEDINFAR SOROLÓGICO

Marketing and Sales of Products for Veterinary Use



in Portugal

# CMO CDMO





# CONTRACT MANUFACTURING

Strict compliance with international requirements and certifications:

## EU GMP

- NP EN ISO 9001:2015
- NP EN ISO 14001:2015
- NP ISO 45001:2019
- EN ISO 22716:2007

**Tech Transfer** - Integrated Project Management Model With Regulatory Support.

## SOLID DOSAGE FORMS

Tablets, Coated Tablets, Pellets, Granules and Capsules

## SEMISOLID DOSAGE FORMS

Creams, Ointments, Gels and Suppositories

## LIQUID DOSAGE FORMS

Syrups, Solutions and Suspensions

INSTALLED CAPACITY:

**130**  
MILLION  
UNITS/YEAR

MORE THAN:

**40**  
CORPORATE  
CUSTOMERS


GROWING BEYOND:

**300**  
QUALIFIED  
PROFESSIONALS

## EU GMP SITE

- FORMULATION DEVELOPMENT
- SCALE-UP
- MANUFACTURING PROCESS DEVELOPMENT & VALIDATION
- ICH STABILITY
- ANALYTICAL METHOD DEVELOPMENT
- ANALYTICAL METHOD VALIDATION

# CONTRACT DEVELOPMENT & LAB SERVICES







# PAIN & FEVER



# PAIN & FEVER



MAIN INGREDIENTS	BRAND	STRENGTH	DOSAGE FORM	PACK	INDICATION/USE
Paracetamol	PARAMOLAN	500 mg	Tablet	20 tabs / 500 tabs	Fever, cold or flu symptoms and pain
Paracetamol	PARACETAMOL GP	1 g	Tablet	20 tabs	Fever, cold or flu symptoms and pain
Paracetamol	PARAMOLAN	125 mg 250 mg	Suppository	5 supp	Fever, cold or flu symptoms and pain
Paracetamol	PARAMOLAN	24 mg/ml	Oral solution	100 ml / 200 ml bottle	Fever, cold or flu symptoms and pain
Paracetamol+Vitamin C	PARAMOLAN C	500 mg + 250 mg	Powder for oral solution	20 sachets	Fever, cold or flu symptoms
Paracetamol+Caffeine	PARAMOLAN CAF	500 mg + 65 mg	Tablet	20 tabs	Fever, cold or flu symptoms and mild to moderate pain

MAIN INGREDIENTS	BRAND	STRENGTH	DOSAGE FORM	PACK	INDICATION/USE
Ibuprofen	TRIFENE 200	200 mg	Film coated tablet	20 tabs / 60 tabs	Mild to moderate pain, fever and cold or flu symptoms
Ibuprofen	TRIFENE Dispersível	200 mg	Dispersible tablet	20 disp tabs	Mild to moderate pain, fever and cold or flu symptoms
Ibuprofen	TRIFENE 400	400 mg	Film coated tablet	20 tabs / 60 tabs	Mild to moderate pain, fever and cold or flu symptoms
Ibuprofen	TRIFENE 600	600 mg	Film coated tablet	20 tabs / 60 tabs	Anti-rheumatic, analgesic, antipyretic
Ibuprofen	TRIFENE	20 mg/ml	Oral suspension	100 ml / 200 ml bottle	Mild to moderate pain and fever



# PARAMOLAN® Caf

Migraine and headache relief



## COUGH & THROAT



MAIN INGREDIENTS	BRAND	STRENGTH	DOSAGE FORM	PACK	INDICATION/USE
Ambroxol, hydrochloride	MUCODRENOL	6 mg/ml	Syrup	200 ml bottle	Mucolytic adjuvant (expectorant)
Bromhexine, hydrochloride	TOSSEQUE	0.8 mg/ml	Syrup	200 ml bottle	Mucolytic adjuvant (expectorant)
Pine buds extract + Eucalyptol + Althea root extract	BEQUISAN	-	Oral Solution	200 ml bottle	Soothing and pleasant effect on throat, pharynx and vocal cords

ESSENTIAL INFORMATION COMPATIBLE WITH THE SUMMARY OF PRODUCT CHARACTERISTICS Name of the medicinal product: Paramolan Caf 500 mg/65 mg tablets Qualitative and quantitative composition: Each tablet contains 500 mg of paracetamol and 65 mg of caffeine. Excipients with known effect: sodium (less than 23 mg per tablet) Pharmaceutical form: Tablet. White oblong tablets. Therapeutic indications: Paramolan Caf is indicated in adults and children aged 12 years and over in the treatment of mild to moderate pain and in febrile states. The situations of mild to moderate pain include: Symptoms associated with flu-like syndromes and colds; Headache, including migraine; Dental pain and pain arising from dental interventions; Odynophagia; Dysmenorrhoea; Muscle aches; Post-traumatic pain. If you do not feel better or if you feel worse after 3 days (in case of fever) or, in case of pain, after 5 days, you must talk to a doctor. Posology and method of administration: Posology: UAdults (including the elderly) and children aged 15 years and over 50 kg: U1-2 tablets, 3-4 times daily (every 6 - 8 hours), as required. Maximum daily dose: 3000 mg/390 mg (paracetamol/caffeine) (6 tablets in 24 hours). U Children 12-15 years (< 50 kg): 1 tablet, 3-4 times daily (every 6 - 8 hours), as required. Maximum daily dose: 2000 mg/260 mg (paracetamol/caffeine) (4 tablets in 24 hours). Maximum daily dose: The dose of 3 g of paracetamol daily, i.e. 6 tablets per day should not be exceeded. An interval of 4 hours (at least) should always be observed between administrations. Paediatric population: Paramolan Caf is not recommended for children under 12 years. Renal impairment: In the event of moderate or severe renal impairment, the dose should be reduced, as follows: Creatinine clearance between 50 - 10 ml/min: 1 tablet, every 6 hours (up to 4 times/day). Creatinine clearance <10 ml/min: 1 tablet, every 8 hours (up to 3 times/day). Hepatic impairment: Patients with mild to moderate hepatic impairment or with Gilbert's syndrome should take this medicine with caution. The dose should be reduced, or the dosing intervals extended. Patients with severe liver impairment should not take this medicinal product. U Method of administration: Oral route. Swallow the tablets with sufficient amount of water. Contraindications: Hypersensitivity to the active substances or to any of the excipients; Severe hepatic impairment; Acute hepatitis; Children under the age of 12 years old. Undesirable effects Adverse reactions have been ranked in frequency using the following convention: very common (≥1/10); common (≥1/100 to <1/10); uncommon (≥1/1,000 to <1/100); rare (≥1/10,000 to <1/1,000); very rare (<1/10,000); not known (cannot be estimated from the available data). Paracetamol: Blood and lymphatic system disorders - Very rare: haematopoiesis disorders (thrombocytopenia, leukopenia, isolated cases of agranulocytosis, pancytopenia); Immune system disorders - Very rare: anaphylaxis and angioedema; Nervous system disorders - Uncommon: vertigo, nervousness; Respiratory, thoracic and mediastinal disorders - Very rare: bronchospasm in predisposed patients (e.g. asthmatic patients sensitive to aspirin and other NSAIDs); Gastrointestinal disorders - Common: nausea, vomiting; Uncommon: diarrhoea, abdominal pain, constipation; Hepatobiliary disorders - Rare: hepatic transaminases increase; Skin and subcutaneous tissue disorders - Rare: erythema; Very rare: cutaneous hypersensitivity reactions, including rash; Not known: toxic epidermal necrolysis and Stevens-Johnson Syndrome. General disorders and administration site conditions disorders - Uncommon: headache, sweating/perspiration, hypothermia; Very rare: allergic reactions, exacerbated hypersensitivity reactions to paracetamol (Quincke's oedema, dyspnoea, sweating, nausea, drop in blood pressure, or even shock). Clinical/epidemiological data seem to indicate that the long-term administration of analgesics may cause nephropathy, including papillary necrosis. Caffeine: The most common undesirable effects of caffeine are nausea caused by irritation of the gastrointestinal tract, as well as insomnia and irritability due to stimulation of the central nervous system. When the recommended dosage of paracetamol-caffeine is combined with dietary intake of caffeine, the resulting dose of caffeine may increase the potential for adverse effects related to caffeine. Psychiatric disorders - Very rare: irritability, insomnia; Gastrointestinal disorders - Very rare: nausea. (11/2021) Product not subject to medical prescription. For more information, please contact the Marketing Authorization Holder. Medinfar Consumer Health - Produtos Farmacêuticos, Lda. Rua Henrique Paiva Couceiro, 27 Venda Nova, 2700-451 Amadora, Portugal





# ANTI- -INFLAMMATORY



# GASTRO- ENTEROLOGY



MAIN INGREDIENTS	BRAND	STRENGTH	DOSAGE FORM	PACK	INDICATION/USE
Piroxicam	REUMOXICAN	20 mg	Suppository Capsule Dispersible tablet	12 supp 20 caps / 60 caps 20 disp tabs / 60 disp tabs	Symptomatic relief of osteoarthritis rheumatoid arthritis and ankylosing spondylitis
Piroxicam	REUMOXICAN Cream	10 mg/g	Cream	100 g tube	Mild to moderate muscular and rheumatic pain
Diethylamine Salicylate	MEDALGINAN Gel	50 mg/g	Gel	40 g tube 100 g tube	Myalgia, lumbago, strains, sprains and muscle spasms
Nimesulide	NIMESULIDA GP	100 mg	Tablet	20 tabs	Acute pain and primary dysmenorrhea
Meloxicam	MELOXICAM GP	7.5 mg 15 mg	Tablet Tablet	20 tabs / 60 tabs 20 tabs / 60 tabs	Symptomatic relief of osteoarthritis, rheumatoid arthritis and ankylosing spondylitis

MAIN INGREDIENTS	BRAND	STRENGTH	DOSAGE FORM	PACK	INDICATION/USE
Omeprazole	PROTON 20 PROTON 40 OMEPRAZOL GP	20 mg 40 mg	GR capsule	14 GR caps 28 GR caps 56 GR caps	GERD and heartburn
Metoclopramide	METOCLOPRAMIDA MEDINFAR	10 mg	Tablet	40 tabs	Prevention and relief of symptoms of nausea and vomiting
Domperidone	CINET	1 mg/ml	Oral suspension	200 ml bottle	Relief of symptoms of nausea and vomiting
Domperidone	CINET DOMPERIDONA GP	10 mg	Tablet	10 tabs / 20 tabs / 60 tabs	Relief of symptoms of nausea and vomiting
Domperidone	CINET	10 mg	Dispersible tablet	20 disp tabs / 60 disp tabs	Relief of symptoms of nausea and vomiting
Phosphate Disodium + Monosodium Phosphate	PRECLINT	240 mg /ml + 542 mg/ml	Oral solution	2 x 45 ml bottle	Osmotic laxative (colonoscopy preparation)
Simethicone	AERO-OM	42 mg	Chewable tablet	60 tabs / 100 tabs	Functional treatment of symptoms associated with gas (abdominal distension, flatulence and meteorism)
Simethicone	AERO-OM	105 mg/ml	Oral emulsion	25 ml bottle	Symptomatic treatment of flatulent colic, difficulties in burping and regurgitation, and functional abdominal distension

# Aero-OM<sup>®</sup>

**FEWER MOMENTS OF COLIC  
MORE MOMENTS AERO-OM<sup>®</sup>**

**GAS | COLIC | BLOATED TUMMY**



## NEWBORNS, BABIES AND CHILDREN

### ESSENTIAL INFORMATION COMPATIBLE WITH THE SUMMARY OF PRODUCT CHARACTERISTICS

Name of the medicinal product: Aero-OM, 105 mg/ml, oral emulsion Qualitative and quantitative composition: One millilitre of oral emulsion contains 105 mg of simethicone as active substance. Excipient with known effect: Sucrose - 300 mg per millilitre of oral emulsion Pharmaceutical form: Oral emulsion. Translucent syrupy white liquid. Therapeutic indications: Symptomatic treatment of flatulent colic, difficulties in burping and regurgitation, and functional abdominal distension. Posology and method of administration: Oral route. Shake before using. Aero-OM oral emulsion: Children aged up to 1 year: 5 to 10 drops 3 or 4 times a day (20 to 40 mg/dose). Children aged 15 years: 10 drops 3 or 4 times a day (40 mg/dose). The duration of treatment should not exceed 10 days. Contraindications: Hypersensitivity to the active substance or to any of the excipients. Undesirable effects: In view of the absence of intestinal absorption, the risk of undesirable effects with simethicone is limited. The incidence of adverse events in published clinical trials is low (1.6%, based on close to 2500 patients exposed), and are primarily gastrointestinal alterations (overall incidence of 1%). The undesirable effects are classified according to the MedDRA convention, by system, organ, class and by frequency as follows: Very common ( $\geq 1/10$ ) Common ( $\geq 1/100$  to  $< 1/10$ ) Uncommon ( $\geq 1/1000$  to  $< 1/100$ ) Rare ( $\geq 1/10,000$  to  $< 1/1000$ ) Very rare ( $< 1/10,000$ ) Unknown (cannot be estimated from the available data). Gastrointestinal alterations: Uncommon: diarrhoea abdominal pain, ileus, nausea and vomiting. Immune system disorders: Rare: hypersensitivity reactions, such as rash, itching, swelling of the face and tongue, and breathing difficulty were also reported. (07/2023) This medicinal product is not subject to medical prescription. For more information, please contact the Marketing Authorization Holder. Medinfar Consumer Health - Produtos Farmacêuticos, Lda. Rua Henrique Paiva Couceiro, 27 Venda Nova, 2700-451 Amadora, Portugal.

## ANTIVIRAL



MAIN INGREDIENTS	BRAND	STRENGTH	DOSAGE FORM	PACK	INDICATION/USE
Aciclovir	CICLOVIRAL	200 mg 400 mg 800 mg	Tablet	25 tabs 25 tabs 25 tabs / 35 tabs / 60 tabs	Herpes simplex and varicella zoster infections
Aciclovir	CICLOVIRAL CREME	50 mg/g	Cream	10 g tube	Cold sores treatment (Herpes simplex infection)

## ANTI-BACTERIAL



MAIN INGREDIENTS	BRAND	STRENGTH	DOSAGE FORM	PACK	INDICATION/USE
Ciprofloxacin, hydrochloride	CIPROFLOXACINA GP	250 mg 500 mg	Tablet	8 tabs / 16 tabs 16 tabs	Bacterial infection



# ciclovir<sup>al</sup> cream

## YOUR SHIELD AGAINST HERPES



### ESSENTIAL INFORMATION COMPATIBLE WITH THE SUMMARY OF PRODUCT CHARACTERISTICS

Name of the medicinal product: CICLOVIRAL 50 mg/g Cream Qualitative and quantitative composition: Each gram of cream contains 50 mg of acyclovir (5% of acyclovir, w/w). Excipients: Each gram of cream contains 67.5 mg cetearyl alcohol; Each gram of cream contains 400 mg propylene glycol. Pharmaceutical form: Cream Therapeutic indications: CICLOVIRAL cream is indicated in the treatment of labial herpes resulting from infection by the Herpes simplex virus type 1. Posology and method of administration: Method of administration: cutaneous use. Adults and children aged 12 or older: CICLOVIRAL cream should be applied five times daily, at approximately 4-hour intervals, omitting the nighttime period. CICLOVIRAL cream should be applied to lesions or imminent lesions, as early as possible after the infection has started. In recurrent episodes, it is particularly important to start treatment in the early stages (prodrome or erythema), or when lesions appear. Studies in patients with labial herpes have demonstrated that treatment is still effective when started after the lesions have appeared (papular and vesicular stages). Treatment should be continued for 4 days. If healing has not occurred after 4 days, treatment may be continued for up to 10 days. If lesions are still present after 10 days, users should be advised to consult a doctor. Children aged 2 to 12 years: The CICLOVIRAL cream doses used in adults are recommended, under medical surveillance, for the following reasons: the stratum corneum is comparable in children and adults; no safety concerns are raised regarding the potential limit of systemic absorption of acyclovir from CICLOVIRAL cream; and there is no evidence to suggest the existence of any differences between age groups regarding the natural history of recurrent episodes of labial herpes. Users should wash their hands before and after applying the cream. Users should also avoid touching or rubbing the lesions, or even touching the lesions with a towel, to avoid aggravating or transferring the infection. Contraindications: Hypersensitivity to acyclovir or any of the excipients. Undesirable effects A transient burning sensation may occur sporadically, following the application of CICLOVIRAL cream. Some cases of mild drying and flaking of the skin have been reported following cutaneous application of acyclovir. Erythema, pruritus and contact dermatitis may occur in rare cases (12/2011) This medicinal product is not subject to medical prescription. For more information, please contact the Marketing Authorization Holder. Medinfar Consumer Health - Produtos Farmacêuticos, Lda. Rua Henrique Paiva Couceiro, 27 Venda Nova, 2700-451 Amadora, Portugal



## CARDIOLOGY



MAIN INGREDIENTS	BRAND	STRENGTH	DOSAGE FORM	PACK	INDICATION/USE
Acetylsalicylic Acid	ASP	100 mg	Tablet	20 tabs / 30 tabs / 60 tabs	Anti-thrombotic
Amiodarone, hydrochloride	AMIODARONA GP	200 mg	Tablet	10 tabs / 60 tabs	Hypertension and angina
Amlodipine, besilate	AMLODIPINA GP	5 mg 10 mg	Tablet	10 tabs / 60 tabs 60 tabs	Tachycardia and arrhythmia associated with several conditions
Captopril	CAPTOPRIL GP	25 mg 50 mg	Tablet	20 tabs / 60 tabs	Hypertension and heart failure
Carvedilol	CARVEDILOL GP	6.25 mg 25 mg	Tablet	14 tabs / 56 tabs	Hypertension, coronary heart disease and heart failure
Indapamide	INDAPAMIDA GP	2.5 mg	Film coated tablet	15 tabs / 60 tabs	Hypertension

# CARDIOLOGY



MAIN INGREDIENTS	BRAND	STRENGTH	DOSAGE FORM	PACK	INDICATION/USE
Lisinopril	LISINOPRIL GP	5 mg 20 mg	Tablet	10 tabs / 60 tabs 60 tabs	Hypertension and heart failure
Losartan, potassium	LOSARTAN GP	50 mg 100 mg	Film coated tablet	14 tabs / 56 tabs	Hypertension and heart failure
Losartan, potassium + Hydrochlorothiazide	LOSARTAN / HCTZ GP	50 mg + 12.5 mg	Film coated tablet	14 tabs / 56 tabs	Hypertension
Ramipril	RAMIPRIL GP	1.25 mg 2.5 mg 5 mg 10mg	Capsule	14 tabs / 56 tabs 56 caps 56 caps 56 caps	Hypertension, heart and renal failure
Ticlopidine, hydrochloride	TICLOPIDINA GP	250 mg	Film coated tablet	10 tabs / 60 tabs	Anti-thrombotic

# CENTRAL NERVOUS SYSTEM



MAIN INGREDIENTS	BRAND	STRENGTH	DOSAGE FORM	PACK	INDICATION/USE
Beta-histine, hydrochloride	BETA-HISTINA GP	16 mg	Tablet	20 tabs / 60 tabs	Meniere's disease
Fluoxetine, hydrochloride	FLUOXETINA GP	20 mg	Capsule	14 tabs / 56 caps	Major depressive episodes, obsessive-compulsive disorder, bulimia nervosa
Paroxetine, hydrochloride	PAROXETINA GP	20 mg	Tablet	10 tabs / 60 tabs	Depression, anxiety disorders



# GLUCOCORTICOID



MAIN INGREDIENTS	BRAND	STRENGTH	DOSAGE FORM	PACK	INDICATION/USE
Deflazacort	DEFLAZACORTE GP	6 mg 30 mg	Tablet	20 / 60 tabs 20 tabs	Anti-inflammatory and immunosuppressant

# CALCIUM FOLINATE



MAIN INGREDIENTS	BRAND	STRENGTH	DOSAGE FORM	PACK	INDICATION/USE
Calcium Folate	MEDIFOLIN	15 mg	Capsule	20 caps	Antidote of folic acid antagonist, megaloblastic anaemia

# WARTS TREATMENT



MAIN INGREDIENTS	BRAND	STRENGTH	DOSAGE FORM	PACK	INDICATION/USE
Salicylic Acid	SALICYLIC ACID	167 mg/g	Solution	10 ml bottle	Warts and callosities treatment

# ANTI-HISTAMINIC



MAIN INGREDIENTS	BRAND	STRENGTH	DOSAGE FORM	PACK	INDICATION/USE
Cetirizine, dihydrochloride	CETIRIZINA GP	10 mg	Film coated tablet	20 tabs	Relief of nasal and ocular symptoms of seasonal and perennial allergic rhinitis, and of chronic idiopathic urticarial

# ANTIDISLIPIDEMIC



MAIN INGREDIENTS	BRAND	STRENGTH	DOSAGE FORM	PACK	INDICATION/USE
Simvastatin	SINVASTINA GP	10 mg 20 mg 40 mg	Film coated tablet	60 tabs 60 tabs 60 tabs	Hypercholesterolemia and cardiovascular prevention

# BENIGN PROSTATIC HYPERTROPHY



MAIN INGREDIENTS	BRAND	STRENGTH	DOSAGE FORM	PACK	INDICATION/USE
Finasteride	FINASTERIDA GP	5 mg	Film coated tablet	60 tabs	Treatment and control of benign prostatic hyperplasia



# VITAMINS & MINERALS



MAIN INGREDIENTS	BRAND	DOSAGE FORM	PACK	INDICATION/USE
Magnesium, Manganese, Copper, Vitamin C, Vitamin B1, Vitamin B2, Vitamin B6	MAGNORANGE ACTIVE	Tablet	30 tabs	Normal energy yielding metabolism, reduction of tiredness and fatigue and normal functioning of the nervous system
Magnesium	MAGNORANGE SENIOR	Oral solution	20 x 10 ml ampoules	Normal energy yielding metabolism, reduction of tiredness and fatigue and maintenance of normal bones
Magnesium, Vitamin C, Ginkgo biloba extract, Vitamin B5, Vitamin B6, Vitamin B1	MAGNORANGE FOCUS	Tablet	60 tabs	Maintenance of memory and good cognitive function, normal physiological function, normal energy yielding metabolism
Echinacea Extract, Vitamin C, Vitamin B6, Zinc, Vitamin D	MAGNORANGE IMUNITAS	Tablet	60 tabs	Normal function of the immune system, normal energy yielding metabolism



# VITAMINS & MINERALS



MAIN INGREDIENTS	BRAND	STRENGTH	DOSAGE FORM	PACK	INDICATION/USE
Magnesium	MAGNORAL	1028.4 mg /10 ml	Oral solution	20 x 10 ml ampoules	Magnesium deficiency and prevention
Vitamin C (Ascorbic Acid)	C'NERGIL	1 g	Powder for oral solution	10 sachets / 20 sachets	Vitamin C deficiency and prevention
Arginine Aspartate + Vitamin B6 + Biotin + Magnesium	BIO RITMO	-	Oral solution	20 x 10 ml ampoules	Normal energy yielding metabolism
Zinc	ZITREX	-	Capsules	60 caps	Protection of cells from oxidative stress and maintenance of normal skin, hair and nails

## AMÉLIORE VOS PERFORMANCES

**1 à 3 ampoules/ jour  
au cours des principaux repas**

**SANS SUCRE**      **SANS ALCOOL**

**ASPARTATE D'ARGININE**      **VITAMINE B6**      **BIOTINE**

**20 AMPOULES EN PLASTIQUE PVC INCASSABLES**  
**UN CONDITIONNEMENT INNOVANT**

Bio-ritmo est un complément alimentaire. Les compléments alimentaires ne doivent pas être utilisés comme substituts d'une alimentation variée. La vitamine B6, la biotine et le magnésium contribuent à un métabolisme énergétique normal ainsi qu'au fonctionnement normal du système nerveux et aussi à des fonctions psychologiques normales. La vitamine B6 et le magnésium contribuent à réduire la fatigue. La vitamine B6 contribue au fonctionnement normal du système immunitaire. La biotine contribue au maintien d'une peau et des cheveux normaux. Le magnésium contribue au maintien d'une assature et d'une dentition normale. Sujets jeunes de plus de 11 ans et les adultes: 1 à 3 ampoules par jour, au cours des principaux repas. Ne pas dépasser les doses journalières recommandées. Les colorants (E 102 et E 124) peuvent réagir de façon croisée avec l'acide acétylsalicylique. Bio-ritmo ne doit pas être pris en cas d'allergie à l'un des ingrédients. Bio-ritmo doit être pris avec précaution chez les patients présentant une insuffisance rénale sévère. Si vous êtes enceinte ou que vous allaitez, demandez l'avis de votre médecin. En cas de doute, consulter la personne responsable de sa mise sur le marché. MEDINFAR CONSUMER HEALTH - PRODUTOS FARMACÉUTICOS, LDA., Rua Henrique Paiva Couceiro, N° 27 Venda Nova, 2700-451 Amadora.





# HAIR LOSS



**PLAYING HIDE AND SEEK WITH YOUR HAIR?**

**WIN EVERY ROUND!**



MAIN INGREDIENTS	BRAND	STRENGTH	DOSAGE FORM	PACK	INDICATION/USE
Minoxidil	TRICOVIVAX	20 mg/ml 50 mg/ml	Cutaneous solution	60 ml / 100 ml bottle (1, 2 and 3 bottles packs)	Androgenic and areata alopecia
Biotin, Niacinamide, Hydrolyzed keratin, Synthetic Peptides	OTÉZIA	-	Lotion	100 ml spray	Anti-hair loss lotion
L-Cystine + Vitamin B6	ONGLINEX	300 mg + 50 mg	Capsule	180 caps	Alopecia, nail dystrophy
L-Cystine, Keratine, Zinc, Iron, Selenium, Niacin, Pantothenic Acid, Vitamin B6, Biotin	TRIZATEK	-	Tablet	60 tablets / 180 tablets	Healthy hair and nails - Hair loss
Caffeine, Niacinamide, Panthenol	TRIKARE C	-	Shampoo	200 ml bottle	Anti-hair loss shampoo

ESSENTIAL INFORMATION COMPATIBLE WITH THE SUMMARY PRODUCT CHARACTERISTICS  
 TRICOVIVAX, 20 mg/ml and 50 mg/ml, cutaneous solution. Qualitative and Quantitative Composition: Tricovivax 20 mg/ml: Each millilitre of cutaneous solution contains 20 mg of minoxidil. Each millilitre of cutaneous solution contains 259 mg of propylene glycol. Tricovivax 50 mg/ml: Each millilitre of topical solution contains 50 mg of minoxidil. Each millilitre of topical solution contains 350 mg of propylene glycol. Complete list of excipients: Propilenoglicol, Etanol a 96%. Pharmaceutical form: Cutaneous Solution. Therapeutic indications: TRICOVIVAX is indicated in the treatment of androgenic alopecia and alopecia areata. Posology and method of administration: Tricovivax is exclusively intended for external use and should only be applied to the scalp. The hair and scalp should be thoroughly dry before topical application of Tricovivax. Do not apply to areas of the body other than the scalp. Tricovivax 20 mg/ml: Apply 1 ml of the 20 mg/ml solution, every 12 hours, pressing the spray pump 6 times. Apply the solution to the center of the affected area and spread it with your fingertips to ensure that the medicinal product is uniformly distributed. For a more localized application, use the applicator provided, which should be fitted onto the top part of the spray head. This dose (1 ml) should be administered regardless of the size of the affected area. The total daily dose should not exceed 2 ml. Tricovivax 50 mg/ml: The recommended dose is 1 ml of the 50 mg/ml solution, every 12 hours. Apply the solution to the center of the affected area and spread it with your fingertips to ensure that the medicinal product is uniformly distributed. This dose (1 ml) should be administered regardless of the size of the affected area. The total daily dose should not exceed 2 ml. Method of application: According to the affected area and the application device provided. Duration of treatment: Studies conducted to the present date indicate that 4 or more months of treatment with Tricovivax may be required for noticeable hair growth. Interruption of treatment may induce a return to the initial condition within 3 to 4 months. Contraindications: Hypersensitivity to the active substance or to any of the excipients. Undesirable effects: Tricovivax 20 mg/ml: Dermatological and sensitivity reactions: The most frequently observed undesirable effects associated with use of minoxidil cutaneous solution are local skin reactions, such as scalp pruritus (1.5% of patients), dry scalp and flaking. Other effects reported were local irritation or burning sensation, including irritative dermatitis. Reported cases of local irritation and allergic contact dermatitis were partly due to the alcohol or propylene glycol present in the formulation. Additionally, wearing of wigs or other types of hairpieces may contribute to local irritation. These effects were generally moderate and only led to treatment discontinuation in rare cases. Allergic reactions including angioedema: frequency not known. Other non-specific allergic and sensitivity reactions have also been reported, including urticaria, allergic rhinitis, facial oedema, eczema, greasy skin, papular rash, folliculitis, local erythema or redness, increased hair loss, alopecia, and hypertrichosis. Cardiovascular disorders: Although not directly attributable to treatment with minoxidil topical solution, cardiovascular effects have been reported, including oedema, precordial pain (usually transient or intermittent), palpitations, increased or decreased blood pressure and/or heart rate, and ECG changes. Nervous system disorders: Although also not attributable to minoxidil, adverse events involving the nervous system have been reported occasionally, including headache (sometimes with exacerbation of pre-existing migraine), dizziness, weakness, taste changes, syncope, "empty head" sensation and vertigo. Cases of anxiety, depression and fatigue have been reported rarely. Other undesirable effects: although also not directly related to topical treatment with minoxidil, have been reported rarely or occasionally, including musculoskeletal disorders (fractures, back pain, retrosternal pain of muscular origin, tendinitis), genitourinary disorders (urinary tract infections, kidney stones, urethritis, prostatitis, epididymitis, impotence), respiratory disorders (upper respiratory tract infections, transient dyspnoea, sinusitis), digestive disorders (diarrhoea, nausea, vomiting, weight increase), blood/immune system disorders (lymphadenopathy, thrombocytopenia), endocrine disorders (breast enlargement), and other disorders (ear infection, conjunctivitis, loss of visual acuity). Tricovivax 50 mg/ml: Tricovivax 50 mg/ml: Common ( $\geq 1/100$ ,  $<1/10$ ); Uncommon ( $\geq 1/1000$ ,  $<1/100$ ); Rare ( $\geq 1/10000$ ,  $<1/1000$ ); Unknown frequency (cannot be estimated from the available data). Skin and subcutaneous tissue disorders: Common: local skin reactions, such as scalp pruritus, dry scalp and flaking. Unknown frequency: local irritation or burning sensation, including irritative dermatitis, eczema, greasy skin, papular rash, folliculitis, local erythema or redness, increased hair loss, alopecia, and hypertrichosis. Reported cases of local irritation and allergic contact dermatitis were partly due to the ethanol or propylene glycol present in the formulation. Additionally, wearing of wigs or other types of hairpieces may contribute to local irritation. Immune system disorders: Unknown frequency: urticaria, allergic rhinitis, facial oedema, allergic reactions including angioedema. The adverse effects listed are not directly attributed to treatment with minoxidil cutaneous solution. Cardiac disorders: Unknown frequency: oedema, precordial pain (usually transient or intermittent), palpitations, increased or decreased blood pressure and/or heart rate, and ECG changes. Nervous system disorders: Uncommon: headache (sometimes with exacerbation of pre-existing migraine), dizziness, weakness, taste changes, syncope, "empty head" sensation and vertigo. Rare: Anxiety, depression, and fatigue. Musculoskeletal and connective tissue disorders: Rare or uncommon: fractures, back pain, retrosternal pain of muscular origin, tendinitis. Genital organs and breast disorders: Rare or uncommon: Prostatitis, epididymitis, and impotence. Renal and urinary disorders: Rare or uncommon: urinary tract infections, kidney stones and urethritis. Respiratory, thoracic and mediastinal disorders: Rare or uncommon: upper respiratory tract infections, transient dyspnoea and sinusitis. Gastrointestinal disorders: Rare or uncommon: diarrhoea, nausea, vomiting, weight increase. Blood and lymphatic system disorders: Rare or uncommon: lymphadenopathy and thrombocytopenia. Endocrine disorders: Rare or uncommon: breast enlargement. Ear and labyrinth disorders: Rare or uncommon: ear infections. Eye disorders: Rare or uncommon: loss of visual acuity and conjunctivitis. These medicinal products are not subject to medical prescription. For more information, please contact the Marketing Authorization Holder. Laboratório Medinfar – Produtos Farmacêuticos, S.A., Rua Henrique de Paiva Couceiro, 29, Venda Nova 2700-451 Amadora. NIF 500 384 045. (07/2019)



# HAIR CARE



MAIN INGREDIENTS	BRAND	DOSAGE FORM	PACK	INDICATION/USE
Panthenol, Hydrolyzed Keratin, Glycolic Acid	PROVIVAX AHA	Shampoo	200 ml tube	Revitalizing and nourishing shampoo
Hydrolyzed Keratin, Biotin, Hamamelis Virginia Leaf Extract	PROVIVAX N	Shampoo	200 ml tube 400 ml bottle	Repairing, nourishing and smoothing shampoo
Glycolic Acid, Biotin, Hydrolyzed keratin, Collagen and Hydrolyzed Wheat Protein	PROVIVAX V	Shampoo	200 ml tube 400 ml bottle	Volumizing, redensifying and revitalizing shampoo



# ANTI-SEBORRHEIC



MAIN INGREDIENTS	BRAND	DOSAGE FORM	PACK	INDICATION/USE
Piroctone Olamine, Ictasol	KETOPAN	Soap	100 g soap	Anti-seborrheic soap
Piroctone Olamine, Zinc PCA, Ictasol	KETOPAN	Foam	200 ml bottle	Anti-seborrheic cleansing foam
Climbazole, Piroctone Olamine, Salicylic Acid and Zinc Lactate	TRIKARE DS	Shampoo	200 ml bottle	Anti-seborrheic and anti-dandruff shampoo





MAIN INGREDIENTS	BRAND	STRENGTH	DOSAGE FORM	PACK	INDICATION/USE
Minocycline	MINOTREX	100 mg	Capsule	16 caps / 32 caps	Skin infections (acne) and other infections
Niacinamide, Glycolic Acid, Salicylic Acid, Oenothera Biennis Oil	NIACIDE	-	Cream	50 g tube	Coadjuvant for acne treatment (dry and sensitive skin)
Niacinamide, Glycolic Acid, Salicylic Acid	NIACIDE	-	Gel	50 g tube	Coadjuvant for acne treatment (oily/combination skin)



MAIN INGREDIENTS	BRAND	STRENGTH	DOSAGE FORM	PACK	INDICATION/USE
Zinc Oxide	HALIBUT	150 mg/g	Ointment	30 g / 50 g/ 100 g tube	Superficial wound healing, nappy rash treatment and first degree burns
Glycerin, Panthenol, Zinc Oxide, Chlorhexidine Digluconate	HALIBUT DERMA	-	Cream	30 g tube	Skin repair for fragile skin and skin discomfort



# DRY & PSORIATIC SKIN



MAIN INGREDIENTS	BRAND	DOSAGE FORM	PACK	USE
Glycerin , Ammonium Lactate, Panthenol, Ceramide NP, Ceramide AP, Ceramide EOP	XERATOP X REPAIR	Lotion	400 ml bottle	Very dry, sensitive and lipid-depleted skin
Paraffinum Liquidum, Ammonium Lactate, Cetyl Alcohol, Panthenol, Cholesterol, Lactic Acid, Ceramide NP, Ceramide AP, Ceramide EOP	XERATOP X REPAIR	Shower Gel	400 ml bottle	Very dry and lipid-depleted skin
Sodium Lactate, Niacinamide, Ceramides, Palmitate, Linoleate, Aloe Barbadensis Leaf Juice, Panthenol	PSORIL MEDINFAR	Lotion	200 ml tube	Coadjuvant in the treatment of skin with psoriasis

**HALIBUT**  
muda fraldas®

Alongside your family ...  
**from day one.**



CLEAN

PROTECTION

REPAIR  
IN CASE OF  
DIAPER RASH

Complete care on diaper change



# BABY CARE



# BABY CARE



MAIN INGREDIENTS	BRAND	DOSAGE FORM	PACK	USE
Helianthus Annuus Seed Oil, Glycerin, Cera Alba, Linum Usitatissimum Seed Oil	HALIBUT muda fraldas	Liniment	200 ml bottle	Cleansing on nappy change
Helianthus Annuus Seed Oil, Triticum Vulgare Bran Extract, Talc, Zinc Oxide	HALIBUT muda fraldas	Cream	100 g / 150 g tube	Nappy rash prevention and skin protection
Zinc Oxide and Miconazole	HALIBUT muda fraldas	Ointment	50 g tube	Skin repair in nappy rash

MAIN INGREDIENTS	BRAND	DOSAGE FORM	PACK	USE
Glycerin, Paraffinum Liquidum, Lanolin	OLEOBAN	Cream	200 g tube / 450 ml bottle	Moisturizing and emollient cream
Helianthus Annuus Seed Oil	OLEOBAN	Bath Solution	300 ml / 500 ml bottle	Moisturizing and emollient bath solution
Helianthus Annuus Seed Oil	OLEOBAN	Soap	100 g Soap	Moisturizing and emollient soap





# BATH & HYDRATION



MAIN INGREDIENTS	BRAND	DOSAGE FORM	PACK	USE
Glycerin, Paraffinum Liquidum, Lanolin	OLEOBAN	Cream	200 g tube / 450 ml bottle / 1 kg bottle	Moisturizing and emollient cream
Ricinus Communis Seed Oil, Panthenol, Helianthus Annuus Seed Oil	OLEOBAN	Shower Gel	300 ml / 500 ml bottle	Moisturizing and emollient shower gel
Helianthus Annuus Seed Oil	OLEOBAN	Soap	100 g soap	Moisturizing and emollient soap
Butyrospermum Parkii Butter, Squalene, Hydrolyzed Linseed Extract	OLEOBAN	Cream	170 ml jar	Intensive moisturizing for very dry skin areas such as heels, elbows, knees and feet; improves skin elasticity and reduces stretch marks
Helianthus Annuus Seed Oil, Ictasol	OLEOBAN DERMA	Bath solution	300 ml bottle	Moisturizing and emollient bath solution for dry and scaly skin; adjuvant in the treatment of skin conditions such as eczema
Polidocanol, Shea Butter, Argania Spinosa Oil	OLEOBAN DERMA	Cream	100 g tube	Hydration of dry, sensitive, scaly and itchy skin



# Oleoban Skin First



Love your skin



# DVINE

DOURO BEAUTY ESSENCE

## ANTI-AGING



MAIN INGREDIENTS	BRAND	DOSAGE FORM	PACK	USE
Albizia Julibrissin Bark Extract, Hyaluronic Acid, Caffeine, Vitis Vinifera Seed Extract, Gold	DVINE GOLD HARVEST Invincible Tawny Eye Serum	Liquid	15 ml bottle	Daily serum for deep anti-aging care of the eye contour
Vitis Vinifera Seed Oil, Ubiquinone, Ectoin, Niacinamide, Caffeine, Gold	DVINE GOLD HARVEST Invincible Tawny Serum	Liquid	50 ml bottle	Serum for premature aging and photo-aging prevention
Vitis Vinifera Seed Oil, Ectoin, Hyaluronic Acid, Phytosterols, Hydrolyzed Pea Protein, Gold	DVINE GOLD HARVEST Invincible Tawny Day Cream	Cream	50 ml bottle	Day cream for intensive revitalization and skin firmness improvement
Vitis Vinifera Seed Extract, Vitis Vinifera Fruit Water, Hyaluronic Acid, Hydrolyzed Pea Protein, Gold	DVINE GOLD HARVEST Invincible Tawny Night Cream	Cream	50 ml bottle	Night cream for a deep regeneration





# ANTI-AGING



MAIN INGREDIENTS	BRAND	DOSAGE FORM	PACK	USE
Vitis Vinifera Fruit Water, Vaccinium Macrocarpon Fruit Powder, Vitis Vinifera Seed Extract	DVINE LIGHT HARVEST Bio Grape Floral Water Cleansing Gel	Gel	125 ml tube	Daily cleansing gel (make-up and impurities removal)
Vitis Vinifera Seed Oil, Vitis Vinifera Fruit Water, Chamomilla Recutita Extract	DVINE LIGHT HARVEST Bio Grape Oil Cleansing Milk	Emulsion	125 ml tube	Daily cleansing milk (make-up and impurities removal)
Jjoba Esters, Vitis Vinifera Seed Powder, Vaccinium Macrocarpon Fruit Powder, Vitis Vinifera Seed Extract	DVINE LIGHT HARVEST Grape Pips Facial Scrub	Liquid	125 ml tube	Facial scrub for impurities, dead cells and excess sebum removal
Vitis Vinifera Fruit Water, Chamomilla Recutita Extract, Epilobium Extract, Cucumis Extract	DVINE LIGHT HARVEST Bio Grape Floral Micellar Water	Solution	200 ml bottle	Cleansing and make up removal of the skin and eyes
Vitis Vinifera Fruit Water, Chamomilla Recutita Extract, Tilia Cordata Extract, Morus Alba Fruit Extract	DVINE LIGHT HARVEST Bio Grape Floral Water Facial Tonic	Solution	200 ml bottle	Facial tonic to remove the impurities, close the pores and fight fatigue



# ANTI-AGING



MAIN INGREDIENTS	BRAND	DOSAGE FORM	PACK	USE
Vitis Vinifera Fruit Water, Albizia Julibrissin Bark Extract, Darutoside, Vitis Vinifera Seed Oil, Hyaluronic Acid	DVINE LIGHT HARVEST Grape Power Dynamic Eye Cream	Cream	15 ml bottle	Daily cream for anti-aging care of the eye contour
Vitis Vinifera Fruit Water, Scenedesmus Rubescens Extract, Hyaluronic Acid, Morus Alba Fruit Extract	DVINE LIGHT HARVEST Grape Power Dynamic Anti-Dark Spot	Liquid	15 ml bottle	Serum for the prevention and correction of the dark-spots
Verbascum Thapsus, Morus Alba Fruit Extract, Vitis Vinifera Seed Oil, Vitis Vinifera Seed Extract	DVINE LIGHT HARVEST Grape Power Dynamic Day Emulsion	Emulsion	50 ml bottle	Day emulsion for anti-aging prevention
Vitis Vinifera Fruit Water, Vitis Vinifera Seed Oil, Verbascum Thapsus, Ectoin, Hyaluronic Acid	DVINE LIGHT HARVEST Grape Power Dynamic Day Cream	Cream	50 ml bottle	Day cream for anti-aging prevention
Prunus Amygdalus Dulcis Oil, Vitis Vinifera Seed Oil, Hyaluronic Acid, Hydrolyzed Pea Protein, Phytosterols	DVINE LIGHT HARVEST Grape Power Dynamic Night Cream	Cream	50 ml bottle	Night cream for anti-aging prevention and skin regeneration

# ANIMAL HEALTH



## DERMA PRODUCTS



MAIN INGREDIENTS	BRAND	STRENGTH	DOSAGE FORM	PACK	INDICATION/USE
Chlorhexidine	HEXADERMA	2%	Shampoo	200 ml bottle	Coadjuvant in the treatment of pyodermatitis and dermatomycosis
Chlorhexidine + Miconazole	HEXADERMA PLUS	2% + 2% 2% + 0,5%	Shampoo Foam	200 ml bottle	Coadjuvant in the treatment of seborrheic dermatitis, superficial pyodermatitis and dermatophytosis
Urea	VETDERMA	10%	Shampoo	200 ml bottle	Coadjuvant in the treatment of dry, peeling and itching skin
Benzoyl Peroxide + Salicylic Acid	SEPIDERMA	25 mg	Shampoo	200 ml bottle	Coadjuvant in the treatment of dry and oil seborrhoea
Phytosphingosine Salicyloyl	ALERDERMA	0,05%	Shampoo	200 ml bottle	Adjuvant in the treatment of pruritus and cutaneous inhalatory conditions in animals with allergic dermatitis, including atopic dermatitis



#### INTERNATIONAL DEPARTMENT

Ana Rita Bivar // [ana.bivar@medinfar.pt](mailto:ana.bivar@medinfar.pt)  
 Cláudia Oliveira // [claudia.oliveira@medinfar.pt](mailto:claudia.oliveira@medinfar.pt)  
 Francisco Costa // [francisco.costa@medinfar.pt](mailto:francisco.costa@medinfar.pt)  
 Manuela Cabugueira // [manuela.cabugueira@medinfar.pt](mailto:manuela.cabugueira@medinfar.pt)  
 Pedro Costa // [pedro.costa@medinfar.pt](mailto:pedro.costa@medinfar.pt)

#### CMO | CDMO

Beatriz Xavier // [beatriz.xavier@medinfar.pt](mailto:beatriz.xavier@medinfar.pt)  
 Paula Leal // [paula.leal@medinfar.pt](mailto:paula.leal@medinfar.pt)  
 Rui Domingos // [rui.domingos@medinfar.pt](mailto:rui.domingos@medinfar.pt)

#### BUSINESS DEVELOPMENT

Francisca Cardoso // [francisca.cardoso@medinfar.pt](mailto:francisca.cardoso@medinfar.pt)  
 Joaquina Couto // [joaquina.couto@medinfar.pt](mailto:joaquina.couto@medinfar.pt)

#### MEDINFAR

Rua Henrique de Paiva Couceiro 29  
 Venda Nova // 2700-451 Amadora // PORTUGAL  
[www.linkedin.com/company/medinfar/](http://www.linkedin.com/company/medinfar/)

#### MEDINFAR MANUFACTURING

Parque Industrial Armando Martins Tavares  
 Rua Outeiro da Armada, 5  
 3150-194 Sebal // PORTUGAL

[www.medinfar.pt](http://www.medinfar.pt)

