



---

**PRODUCT LIST**  
FOR EXPORT

---



## Contents

### Artificial Tears 4p~5p



Lacure  
Ophthalmic Solution



Optagent  
Ophthalmic Solution



EYE:O mini  
Ophthalmic Solution



Reva-K  
Ophthalmic Solution

### Antibiotics 5p~7p



OcuFloX  
Ophthalmic Solution



OcuFloX  
Ophthalmic Ointment



OcuLevo  
Ophthalmic Solution



OcuRacin  
Ophthalmic Solution



OcuRacin  
Ophthalmic Ointment



Cifex  
Ophthalmic Solution



Eymox  
Ophthalmic Solution

### Antibiotics + Steroids 7p~8p



Forus  
Ophthalmic Solution



Forus  
Ophthalmic Ointment

### Antiallergic 8p~9p



Allerfen  
Ophthalmic Solution



Allinol  
Ophthalmic Solution



OcuOlo  
Ophthalmic Solution

**NSAIDs**  
9p



Optanac  
Ophthalmic Solution

**Miotics**  
9p



OcuCarpine  
Ophthalmic Solution

**Mydriatics**  
10p



OcuHomapine  
Ophthalmic Solution



OcuCyclo  
Ophthalmic Solution

**Steroids**  
10p



OcuMetholone  
Ophthalmic Solution 0.02% / 0.1%

**Antivirus**  
11p



OcuFridine  
Ophthalmic Solution



Herpecid  
Ophthalmic Ointment

**Cataracts**  
11p



Curelen  
Ophthalmic Solution

Artificial Tears

**Lacure**  
Ophthalmic Solution



Indication	Keratoconjunctivitis sicca with following symptoms <b>Endogenous disease:</b> Sjögren's syndrome, Stevens-Johnson syndrome, dry eye syndrome <b>Enthetic disease:</b> Drug-induced trauma after operation, enthetic disease due to the contact lenses
Active Ingredient	Sodium Hyaluronate 1 mg/mL
Description	Clear, colorless aqueous ophthalmic solution with viscosity
Dosage & Administration	Instill 1 drop into the affected eye(s) 5~6 times daily
Package	5 mL/bottle, 10 mL/bottle

Artificial Tears

**Optagent**  
Ophthalmic Solution



Indication	Symptomatic relief of irritated dry eyes and symptoms of dry eyes
Active Ingredient	Povidone 20 mg/mL
Description	Clear, colorless aqueous ophthalmic solution
Dosage & Administration	Instill 1~2 drop(s) into the affected eye(s) 4~5 times daily
Package	10 mL/bottle

Artificial Tears

**EYE2O mini**  
Ophthalmic Solution

PRESERVATIVE FREE



Indication	Lubricant to relieve irritation and discomfort due to dryness of the eye or due to exposure to wind or sun
Active Ingredient	Carboxymethylcellulose Sodium 5 mg/mL
Description	Clear, colorless to pale yellow ophthalmic solution in a single-use container
Dosage & Administration	Instill 1~2 drop(s) into affected eye(s), and discard the remaining liquid and container after instillation
Package	0.5 mL/tube (30 tubes/package)

Artificial Tears

**Reva-K**  
Ophthalmic Solution



Indication	Improvement of corneal and conjunctival epithelial damage in adult patients with dry eyes
Active Ingredient	Rebamipide 20 mg/mL
Description	Clear, colorless liquid in a translucent plastic container
Dosage & Administration	Instill 1 drop into affected eye(s) 4 times daily.
Package	5 mL/bottle

Antibiotics

**OcuFlox**  
Ophthalmic Solution

PRESERVATIVE FREE



Indication	Topical treatment of blepharitis, dacryocystitis, hordeolum, conjunctivitis, tarsadenitis, and keratitis (including corneal ulcer) and also as an aseptic measure in ophthalmic surgeries pre- and postoperatively
Active Ingredient	Ofloxacin 3 mg/mL
Description	Clear, pale yellow to lemon-yellow ophthalmic solution
Dosage & Administration	Instill 1 drop into the affected eye(s) 3 times daily <b>Post-operative infections:</b> Instill 5 times daily for 2 days pre-operatively, make adjustments properly for the application of this medication on the day of operation based upon the time and the type of the surgeries, and instill the medication each time when changing the bandage afterwards
Package	5 mL/bottle

Antibiotics

**OcuFlox**  
Ophthalmic Ointment

PRESERVATIVE FREE



Indication	Topical treatment of blepharitis, dacryocystitis, hordeolum, conjunctivitis, tarsadenitis, and keratitis (including corneal ulcer) and also as an aseptic measure in ophthalmic surgeries pre- and postoperatively
Active Ingredient	Ofloxacin 3 mg/g
Description	Odorless, light yellow ophthalmic ointment
Dosage & Administration	Apply a small amount of OcuFlox into the affected eye(s) 3 times daily <b>Post-operative infections:</b> Apply 5 times daily for 2 days pre-operatively, make adjustments properly for the application of this medication on the day of operation based upon the time and the type of the surgeries, and apply the medication each time when changing the bandage afterwards
Package	5 g/tube

Antibiotics

**OcuLevo**  
**Ophthalmic Solution 0.5%**

**PRESERVATIVE FREE**



Indication	Topical treatment of ocular bacterial infections including blepharitis, dacryocystitis, hordeolum, conjunctivitis, tarsadenitis, keratitis, corneal ulcer, and also as an aseptic measure in ophthalmic surgeries
Active Ingredient	Levofloxacin Hydrate 5 mg/mL
Description	Clear, pale yellow to lemon-yellow aqueous ophthalmic solution
Dosage & Administration	Instill 1 drop into the affected eye(s) 3 times daily
Package	5 mL/bottle

Antibiotics

**OcuRacin**  
**Ophthalmic Solution**



Indication	Topical treatment of ocular bacterial infections including blepharitis, dacryocystitis, hordeolum, conjunctivitis, keratitis, and corneal ulcer
Active Ingredient	Tobramycin 3 mg/mL
Description	Colorless to pale yellow aqueous ophthalmic solution
Dosage & Administration	<b>Mild to moderate infection:</b> Instill 1-2 drop(s) into the affected eye(s) every 4 hours. <b>Severe infection:</b> Instill 2 drops into the affected eye(s) hourly until the conditions get improved gradually decreasing the dose to discontinuation
Package	5 mL/bottle

Antibiotics

**OcuRacin**  
**Ophthalmic Ointment**



Indication	Topical treatment of ocular bacterial infections including blepharitis, dacryocystitis, hordeolum, conjunctivitis, keratitis, and corneal ulcer
Active Ingredient	Tobramycin 3 mg/g
Description	White or pale yellow ointment
Dosage & Administration	<b>Mild to moderate infection:</b> Apply a small amount (about 1 cm) into the affected eye(s) 2-3 times daily <b>Severe infection:</b> Apply a small amount (about 1 cm) into the affected eye(s) every 3-4 hours until the conditions get improved gradually decreasing the dose to discontinuation
Package	5 g/tube

Antibiotics

## Cifex Ophthalmic Solution



<b>Indication</b>	<b>Corneal ulcers:</b> <i>Pseudomonas aeruginosa</i> , <i>Serratia marcescens</i> , <i>Staphylococcus aureus</i> , <i>Streptococcus pneumoniae</i> , <i>Staphylococcus epidermidis</i> <b>Bacterial conjunctivitis:</b> <i>Staphylococcus aureus</i> , <i>Staphylococcus epidermidis</i> , <i>Streptococcus pneumoniae</i>
<b>Active Ingredient</b>	Ciprofloxacin Hydrochloride 3.5 mg/mL
<b>Description</b>	Clear, colorless aqueous ophthalmic solution
<b>Dosage &amp; Administration</b>	<b>Corneal ulcers:</b> (1) On the first day, instill 2 drops into the affected eye(s) every 15 minutes for the first 6 hours and then 2 drops every 30 minutes for the remainder of the day (2) On the second day, instill 2 drops in the affected eye(s) hourly. (3) On the third day through the fourteenth day, place 2 eye drops in the affected eye(s) every 4 hours (4) Treatment may be continued after 14 days if corneal re-epithelialization has not occurred <b>Bacterial conjunctivitis:</b> Instill 2 drops into the affected eye(s) every 2 hours while awake for the first 2 days and 1~2 drop(s) every 4 hours while awake for the next 5 days
<b>Package</b>	5 mL/bottle

Antibiotics

## Eymox Ophthalmic Solution

PRESERVATIVE FREE



<b>Indication</b>	Bacterial conjunctivitis, tarsadenitis, keratitis, corneal ulcer, and also as an aseptic measure in ophthalmic surgeries
<b>Effective Bacillus Strain</b>	<i>Staphylococcus</i> , <i>Streptococcus</i> , <i>Streptococcus pneumonia</i> , <i>Micrococcus</i> , <i>Enterococci</i> , <i>Corynebacterium spp.</i> , <i>Citrobacter spp.</i> , <i>Klebsiella spp.</i> , <i>Enterobacter spp.</i> , <i>Serratia spp.</i> , <i>Proteus spp.</i> , <i>Morganella morganii</i> , <i>influenza bacteria</i> , <i>Pseudomonas aeruginosa</i> , <i>Burkholderia cephalcy</i> , <i>Stenotropomonas (Santomonas) Maltophilia</i> , <i>Acinetobacter spp.</i> , <i>Propionibacterium acnes</i>
<b>Active Ingredient</b>	Moxifloxacin Hydrochloride 5.45 mg/mL
<b>Description</b>	Light greenish yellow ophthalmic solution
<b>Dosage &amp; Administration</b>	<b>Bacterial conjunctivitis, tarsadenitis, keratitis, corneal ulcer:</b> Instill 1 drop into the affected eye(s) 3 times daily <b>Pre-operation:</b> Instill 1 drop into the affected eye(s) 5 times daily <b>Post-operation:</b> Instill 1 drop into the affected eye(s) 3 times daily
<b>Package</b>	5 mL/bottle

Antibiotics+Steroids

## Forus Ophthalmic Solution



<b>Indication</b>	Treatment of conjunctivitis, blepharitis, and keratitis caused by strains of the microorganisms susceptible to polymyxin B sulfate and neomycin sulfate
<b>Active Ingredient</b>	Neomycin Sulfate 3.5 mg/mL (potency) Dexamethasone 1 mg/mL Polymyxin B Sulfate 6,000 IU/mL
<b>Description</b>	White ophthalmic solution
<b>Dosage &amp; Administration</b>	Instill 1~2 drop(s) into the conjunctival sac(s) of the affected eye(s) 4~6 times daily. In severe infections, instill the drug hourly until the inflammation subsides being tapered to discontinuation.
<b>Package</b>	5 mL/bottle



Antibiotics+Steroids

**Forus**  
**Ophthalmic Ointment**

PRESERVATIVE FREE



<b>Indication</b>	Treatment of conjunctivitis, blepharitis, and keratitis caused by strains of the microorganisms susceptible to polymyxin B sulfate and neomycin sulfate
<b>Active Ingredient</b>	Neomycin Sulfate 3.5 mg/g (potency) Dexamethasone 1 mg/g Polymyxin B Sulfate 6,000 IU/g
<b>Description</b>	White to pale yellow ophthalmic ointment
<b>Dosage &amp; Administration</b>	Apply small amount into the conjunctival sac(s) 3~4 times daily or use concomitantly with ophthalmic solution at bedtime
<b>Package</b>	5 g/tube

Antiallergic

**Allerfen**  
**Ophthalmic Solution**



<b>Indication</b>	Allergic conjunctivitis
<b>Active Ingredient</b>	Ketotifen Fumarate 0.69 mg/mL
<b>Description</b>	Clear, colorless to pale yellow ophthalmic solution
<b>Dosage &amp; Administration</b>	<b>Adults:</b> Instill 1 drop into the affected eye(s) 2~4 times daily <b>Infants/children:</b> As prescribed by the physician
<b>Package</b>	5 mL/bottle

Antiallergic

**Allinol**  
**Ophthalmic Solution**



<b>Indication</b>	Treatment of eye itching associated with allergic conjunctivitis
<b>Active Ingredient</b>	Olopatadine Hydrochloride 7.76mg/mL
<b>Description</b>	Colorless to light yellow liquid in an opaque white plastic container
<b>Dosage &amp; Administration</b>	Instill 1 drop into affected eye two times daily
<b>Package</b>	5 mL/bottle



Antiallergic

**OcuOlo**  
Ophthalmic Solution



Indication	Treatment of symptoms due to allergic conjunctivitis
Active Ingredient	Olopatadine Hydrochloride 7.76mg/mL
Description	Colorless to light yellow liquid in an opaque white plastic container
Dosage & Administration	Instill 1 drop into affected eye two times daily
Package	2.5 mL/bottle

NSAIDs

**Optanac**  
Ophthalmic Solution



Indication	<b>Prevention of the following symptoms in cataract surgeries:</b> Post-operative inflammation, intraoperative miosis, cystoid macular edema related with lens extraction and intraocular lens implantation <b>Treatment of inflammation in anterior segment of the eyes:</b> Non-infectious inflammations including chronic conjunctivitis and post-traumatic inflammation
Active Ingredient	Diclofenac Sodium 1 mg/mL
Description	Colorless or pale yellow, clear, aqueous ophthalmic solution
Dosage & Administration	<b>For cataract surgery:</b> 1) Preoperative: Instill 1 drop into the affected eye(s) 5 times during 3 hours of time 2) Right after the surgery: Instill 1 drop into the affected eye(s) 3 times 3) Maintenance: Instill 1 drop into the affected eye(s) 3-5 times daily (administration period can be adjusted based upon the symptoms)  <b>For inflammation in the anterior segment of the eyes:</b> Instill 1 drop into the affected eye(s) 4-5 times daily
Package	5 mL/bottle

Miotics

**OcuCarpine**  
Ophthalmic Solution 2%



Indication	Treatment of glaucoma and miotic for diagnosis or treatment
Active Ingredient	Pilocarpine Hydrochloride 20 mg/mL
Description	Colorless aqueous ophthalmic solution
Dosage & Administration	Instill 1~2 drop(s) into the affected eye(s) every 1~3 times daily
Package	10 mL/bottle

Mydriatics

**OcuHomapine**  
Ophthalmic Solution



Indication	Mydriasis and cycloplegia for diagnosis and treatment
Active Ingredient	Homatropine Hydrobromide 20 mg/mL
Description	Colorless aqueous ophthalmic solution
Dosage & Administration	Instill 1~2 drop(s) into the affected eye(s) 1 time a day
Package	5 mL/bottle

Mydriatics

**OcuCyclo**  
Ophthalmic Solution



Indication	Mydriasis and cycloplegia for diagnosis and treatment
Active Ingredient	Cyclopentolate Hydrochloride 10 mg/mL
Description	Colorless aqueous ophthalmic solution
Dosage & Administration	Instill 1 drop into the affected eye(s) every 1~3 times daily or instill 1 drop additionally 5~10 minutes later
Package	5 mL/bottle

Steroids

**OcuMetholone**  
Ophthalmic Solution  
0.02% / 0.1%



Indication	<b>For 0.02%:</b> Inflammatory diseases of the external ocular segments (including blepharitis, conjunctivitis, keratitis, scleritis, episcleritis, etc.) <b>For 0.1%:</b> Inflammatory diseases of the external and anterior ocular segments (including blepharitis, conjunctivitis, keratitis, scleritis, episcleritis, iritis, iridocyclitis, uveitis, and postoperative inflammation, etc.)
Active Ingredient	Fluorometholone 0.2 mg/mL, 1 mg/mL
Description	Pale milky white, aqueous suspension
Dosage & Administration	Instill 1~2 drop(s) into the affected eye(s) 2~4 times daily
Package	5 mL/bottle

Antivirus

**OcuFridine**  
Ophthalmic Solution



Indication	Treatment of herpetic keratitis
Active Ingredient	Trifluridine 10 mg/mL
Description	Clear, colorless aqueous ophthalmic solution
Dosage & Administration	Instill 1 drop into the affected eye(s) every 2 hours daily Instill 1 drop 5~7 times daily until healed and 1 drop 2~3 times daily for additional 7 days
Package	5 mL/bottle

Antivirus

**Herpecid**  
Ophthalmic Ointment



Indication	Treatment of keratitis caused by herpes simplex virus
Active Ingredient	Acyclovir 30 mg/g
Description	White to pale yellow ophthalmic ointment
Dosage & Administration	The usual dose is a 1 cm ribbon of ointment 5 times a day at approximately 4 hourly intervals Apply a 1 cm ribbon of ointment to the inside of the lower conjunctival sac. Treatment should continue for at least 3 days after healing is completed
Package	3.5 g/tube

Cataracts

**Curelen**  
Ophthalmic Solution



Indication	Cloudy and hemorrhage of hyaline bodies due to aging, myopia, hypertension, diabetes mellitus, periphlebitis, initial symptom of senile cataract
Active Ingredient	Potassium Iodide 3 mg/ml Sodium Iodide 3 mg/ml
Description	Colorless ophthalmic solution
Dosage & Administration	Instill 1 drop into the affected eye(s) every 1~3 times daily
Package	10 ml/bottle



MEMO



A large area of the page containing multiple horizontal dotted lines, intended for writing a memo.



## PRODUCT LIST

Category	No.	Product	Package
Artificial Tears	1	Lacure Ophthalmic Solution	5 mL/bottle, 10 mL/bottle
	2	Optagent Ophthalmic Solution	10 mL/bottle
	3	EYE:O mini Ophthalmic Solution	0.5 mL/tube (30 tubes/package)
	4	Reva-K Ophthalmic Solution	5 mL/bottle
Antibiotics	5	OcuFlox Ophthalmic Solution	5 mL/bottle
	6	OcuFlox Ophthalmic Ointment	5 g/tube
	7	OcuLevo Ophthalmic Solution 0.5%	5 mL/bottle
	8	OcuRacin Ophthalmic Solution	5 mL/bottle
	9	OcuRacin Ophthalmic Ointment	5 g/tube
	10	Cifex Ophthalmic Solution	5 mL/bottle
	11	Eymox Ophthalmic Solution	5 mL/bottle
Antibiotics + Steroids	12	Forus Ophthalmic Solution	5 mL/bottle
	13	Forus Ophthalmic Ointment	5 g/tube
Antiallergic	14	Allerfen Ophthalmic Solution	5 mL/bottle
	15	Allinol Ophthalmic Solution	5 mL/bottle
	16	OcuOlo Ophthalmic Solution	2.5 mL/bottle
NSAIDs	17	Optanac Ophthalmic Solution	5 mL/bottle
Miotics	18	OcuCarpine Ophthalmic Solution 2%	10 mL/bottle
Mydriatics	19	OcuHomapine Ophthalmic Solution	5 mL/bottle
	20	OcuCyclo Ophthalmic Solution	5 mL/bottle
Steroids	21	OcuMetholone Ophthalmic Solution 0.02% / 0.1%	5 mL/bottle
Antivirus	22	OcuFridine Ophthalmic Solution	5 mL/bottle
	23	Herpecid Ophthalmic Ointment	3.5 g/tube
Cataracts	24	Curelen Ophthalmic Solution	10 ml/bottle





---

## PRODUCT LIST

---

*Samil*

---

[KOREA]

**Head Office**

(06666) 155 Hyoryeong-ro, Seocho-gu, Seoul,  
Republic of Korea  
TEL +82-2-520-0300 FAX +82-2-588-5924

**Ansan Plant**

(15433) 216 Sandan-ro, Danwon-gu, Ansan-si,  
Gyeonggi-do, Republic of Korea  
TEL +82-31-490-4000 FAX +82-31-492-3134

**Samil Eyecare Innovation Center**

3rd floor, 222 Banpo-Daero, Seocho-Gu, Seoul,  
Republic of Korea  
TEL +82-2-520-0300 FAX +82-2-588-5924

[VIETNAM]

**Vietnam Representative Office**

18th Floor, CJ Building, 2Bis4-6 Le Thanh Ton,  
Ben Nghe Ward, District 1, HCMC, Vietnam  
TEL +84-28-6255-6890 FAX +84-28-6255-6801

**S1 Plant**

Lot-15, Street D15, Hi-tech Industrial Park,  
Tnag Nhon Phu B Ward, District 9 Ho Chi Minh City, Vietnam  
TEL +84-28-8887-3033

[CANADA]

**Samil Pharmaceutical North America**

1055 West Georgia Street 1500 Royal Centre, P.O. Box 11117  
Vancouver, BC V6E 4N7, Canada  
TEL +1-778-372-8370

**Canada Liaison Office**

(Office 903) 900-2025 Willingdon Ave.,  
Burnaby, BC V5C 0J3, Canada  
TEL +1-778-372-8370

---